


## Solutions to measure for depressive symptoms across the age span

### Screen children for depressive symptoms

Designed to measure depression in children ages 7 to 13 years, the **Reynolds Child Depression Scale™—2nd Edition (RCDS™-2)** is a 30-item self-report form. Children are asked to answer questions about how they have been feeling during the past 2 weeks. Administration time is 15 minutes. Available in paper and pencil or **online via PARiConnect**. The RCDS-2: Short Form (RCDS-2™:SF), ideal for screening, is also available and takes just 2 to 3 minutes to administer.


Learn more at [parinc.com/RCDS2](http://parinc.com/RCDS2) or [parinc.com/RCDS2\\_SF](http://parinc.com/RCDS2_SF)




### Measure the severity of depressive symptoms in adolescents

The **Reynolds Adolescent Depression Scale, 2nd Edition (RADS-2™)**, is a 30-item self-report form that measures four basic dimensions of depression in adolescents ages 11 to 20 years. Standard scores and associated clinical cutoff score provide an indication of the clinical severity of the individual's depressive symptoms. Administration time is 10 minutes. Available in paper and pencil or **online via PARiConnect**. The RADS-2: Short Form (RADS-2™:SF), ideal for screening, is also available and takes just 2 to 3 minutes to administer.

Learn more at [parinc.com/RADS2](http://parinc.com/RADS2) or [parinc.com/RADS2\\_SF](http://parinc.com/RADS2_SF)





## Measure depressive symptomatology across the age range


The **Clinical Assessment of Depression™ (CAD™)** is a self-report measure that is sensitive to depressive symptomatology throughout the life span (ages 8 to 79 years). Administration takes 10 minutes. Available in paper and pencil, **online via PARiConnect**, and software.

Learn more at [parinc.com/CAD](http://parinc.com/CAD)

## Screen for depression in adults

The **Reynolds Depression Screening Inventory™ (RDSI™)** measures the severity of contemporary depressive symptoms in adults ages 18 to 89 years. Although a formal diagnosis is not provided, empirically derived cutoff scores identify individuals who may be at risk for more serious, diagnostic forms of depression. Administration time is 5 to 10 minutes. Available in paper and pencil.

Learn more at [parinc.com/RDSI](http://parinc.com/RDSI)



The **Patient Health Questionnaire-9 (PHQ-9)** is a nine-question instrument used in the screening and detection of depression symptoms. Administration time is 5 to 10 minutes. Available online and via CheckIT on PARiConnect.

Learn more at [parinc.com/checkit](http://parinc.com/checkit)

## PUT OUR EXPERIENCE TO WORK FOR YOU!

Learn more from  
your **PAR Healthcare**  
Clinical Assessment Advisor



**Melissa Milanak, PhD**  
813.955.2927 or  
[mmilanak@parinc.com](mailto:mmilanak@parinc.com)

Learn more at [parinc.com](http://parinc.com)